arXiv:1604.07116v1 [math.CV] 25 Apr 2016

ANALYTIC ADJOINT IDEAL SHEAVES ASSOCIATED TO PLURISUBHARMONIC FUNCTIONS

QI'AN GUAN AND ZHENQIAN LI

Abstract. In this article, we will present that the analytic adjoint ideal sheaves associated to plurisubharmonic functions are not coherent.

1. Introduction

The adjoint ideal sheaf on a smooth complex algebraic variety X is a variant of the multiplier ideal sheaf in algebraic geometry (see [1, 3, 4] for more details).

In [1], Guenancia gave an analytic definition of an adjoint ideal sheaf associated to a quasi-plurisubharmonic function φ along a simple normal crossing (SNC) divisor $D=\sum D_i$ and established the compatibility with the algebraic adjoint ideal whenever φ has analytic singularities.

Let X be a complex manifold, $D=\sum D_i$ an SNC divisor and φ a quasi-plurisubharmonic function on X. Let $Adj^{\alpha}_{D,*}(\varphi)\subset\mathcal{O}_X$ be the ideal sheaf of germs of holomorphic functions $f\in\mathcal{O}_{X,x}$ such that

$$|f|^2\prod_{k=1}^{p}\frac{1}{|h_k|^2(-\log|h_k|)^{\alpha}}e^{-\varphi}$$

is integrable with respect to the Lebesgue measure in some local coordinates near x, where $h=h_1\cdots h_p$ is the minimal defining function of D near x and $\alpha>1$.

In [1] (see also [2]), Guenancia gave the following analytic definition of adjoint ideal sheaf, which generalized the algebraic adjoint ideal sheaf (Proposition 2.11 in [1]; see also Proposition 5.1 in [2]).

Definition 1.1. ([1, 2]). The ideal sheaf $Adj^{\alpha}_{D}(\varphi):=\cup_{\varepsilon>0}Adj^{\alpha}_{D,*}((1+\varepsilon)\varphi)$ is called the *analytic adjoint ideal sheaf* associated to φ along D.

Note that in [1] Guenancia used $\alpha=2$ in the definition, and later Kim in [2] extended the definition to $\alpha>1$ case. When e^{φ} is locally Hölder continuous, Guenancia established the coherence of Adj^{α}_{D} for smooth divisor D with $\varphi|_D\not\equiv-\infty$ (see Corollary 2.19 in [1]).

As mentioned by Guenancia and Kim, it is natural to ask the following

Question 1.1. ([1, 2]). For $\alpha>1$ and φ a general quasi-plurisubharmonic function on X, is the analytic adjoint ideal sheaf $Adj^{\alpha}_{D}(\varphi)$ coherent?

In this article, we will present the following negative answer to Question 1.1.

Date: April 26, 2016.
2010 *Mathematics Subject Classification.* 32C25, 32C35, 32U05.
The first author was partially supported by NSFC-11522101 and NSFC-11431013.
Key words. Plurisubharmonic function, Adjoint ideal sheaf, Hölder continuous.

Theorem 1.1. *There exists a plurisubharmonic function φ and a smooth divisor D with $\varphi|_D \not\equiv -\infty$ near the origin $o \in \mathbb{C}^n$ $(n \geq 3)$ such that for any $\alpha > 1$, the analytic adjoint ideal sheaf $Adj_D^\alpha(\varphi)$ is not coherent near o.*

Specifically, we will construct φ and D with $\varphi|_D \not\equiv -\infty$ near o such that the zero set of $Adj_D^\alpha(\varphi)$ is not an analytic set near o.

2. Proof of main results

We are now in a position to prove Theorem 1.1.

***Proof of Theorem* 1.1.** Let $D = \{z_1 = 0\}$ and

$$\varphi(z) = \max\{\sum_{k=2}^{\infty} \alpha_k \log(|z_1| + |z_2 - \frac{1}{k}|^{\beta_k}), \lambda \log|z_3|\},$$

where $\alpha_k = \frac{1}{2^{k!}}$, $\beta_k = 3 \cdot 2^{k!}$ and $\lambda > 6$. Then D is smooth and $\varphi(z)$ is a plurisubharmonic function near $o \in \mathbb{C}^n$ $(n \geq 3)$. Without loss of generality, we assume $n = 3$ and U contained in the unit polydisk Δ^3 is a neighborhood of o such that $\log(|z_1| + |z_2 - \frac{1}{k}|^{\beta_k}) < 0$ for any $k \geq 2$ on U.

Step 1. *Nonintegrability of* $\frac{1}{|z_1|^2(-\log|z_1|)^\alpha} e^{-\varphi}$ *near* $(0, \frac{1}{k}, 0)$ *for any* $(0, \frac{1}{k}, 0) \in U$. Since

$$\begin{aligned}
\varphi(z) &= \max\{\sum_{k=2}^{\infty} \alpha_k \log(|z_1| + |z_2 - \frac{1}{k}|^{\beta_k}), \lambda \log|z_3|\} \\
&\leq \max\{\alpha_k \log(|z_1| + |z_2 - \frac{1}{k}|^{\beta_k}), \lambda \log|z_3|\} \\
&\leq \log\big((|z_1| + |z_2 - \frac{1}{k}|^{\beta_k})^{\alpha_k} + |z_3|^\lambda\big)
\end{aligned}$$

near o, replacing $z_2 - \frac{1}{k}$ by z_2 near $(0, \frac{1}{k}, 0)$, for sufficiently small polydisk Δ_r^3 we have

$$\begin{aligned}
&\int_{\Delta_r^3} \frac{1}{|z_1|^2(-\log|z_1|)^\alpha} e^{-\varphi(z_1, z_2 + \frac{1}{k}, z_3)} dV_3 \\
\geq &\int_{\Delta_r^3} \frac{dV_3}{\big((|z_1| + |z_2|^{\beta_k})^{\alpha_k} + |z_3|^\lambda\big)|z_1|^2(-\log|z_1|)^\alpha} \\
= &\int_{\Delta_r^*} \frac{dV_1}{|z_1|^2(-\log|z_1|)^\alpha} \int_{\Delta_r^2} \frac{|z_1|^{\frac{2}{\beta_k} + \frac{2\alpha_k}{\lambda}}}{|z_1|^{\alpha_k}} \frac{(\frac{\sqrt{-1}}{2})^2 d(\frac{z_2}{|z_1|^{\frac{1}{\beta_k}}}) \wedge d(\frac{\bar{z}_2}{|z_1|^{\frac{1}{\beta_k}}}) \wedge d(\frac{z_3}{|z_1|^{\frac{\alpha_k}{\lambda}}}) \wedge d(\frac{\bar{z}_3}{|z_1|^{\frac{\alpha_k}{\lambda}}})}{(1 + \frac{|z_2|^{\beta_k}}{|z_1|})^{\alpha_k} + \frac{|z_3|^\lambda}{|z_1|^{\alpha_k}}} \\
\geq &\int_{\Delta_r^*} \frac{dV_1}{|z_1|^{2 + (\alpha_k - \frac{2}{\beta_k} - \frac{2\alpha_k}{\lambda})}(-\log|z_1|)^\alpha} \int_{\Delta} \frac{(\frac{\sqrt{-1}}{2})^2 dw_2 \wedge d\bar{w}_2 \wedge dw_3 \wedge d\bar{w}_3}{(1 + |w_2|^{\beta_k})^{\alpha_k} + |w_3|^\lambda} \\
\geq C \cdot &\int_{\Delta_r^*} \frac{dV_1}{|z_1|^{2 + (\alpha_k - \frac{2}{\beta_k} - \frac{2\alpha_k}{\lambda})}(-\log|z_1|)^\alpha} = +\infty,
\end{aligned}$$

where $C > 0$ is some constant and the last equality is from $\alpha_k - \frac{2}{\beta_k} - \frac{2\alpha_k}{\lambda} > 0$. It follows that $\frac{1}{|z_1|^2(-\log|z_1|)^\alpha} e^{-\varphi}$ is not locally integrable near $(0, \frac{1}{k}, 0)$ for any $(0, \frac{1}{k}, 0) \in U$.

Step 2. *Integrability of* $\frac{1}{|z_1|^2(-\log|z_1|)^\alpha}e^{-(1+\varepsilon)\varphi}$ *near* $(0,z_2,0)\in U$ *with* $z_2\neq\frac{1}{k},0$.

Take $0<\varepsilon_0<1$ such that $\alpha-\varepsilon_0>1$ and $|z_2-\frac{1}{k}|>\varepsilon_0$ for any k. Then, for any $N\geq 1$ with $\varepsilon_0^{\beta_{N+1}}<|z_1|\leq\varepsilon_0^{\beta_N}$, we obtain

$$\begin{aligned}\varphi(z)&\geq\sum_{k=2}^{\infty}\alpha_k\log(|z_1|+|z_2-\frac{1}{k}|^{\beta_k})\geq\sum_{k=2}^{\infty}\alpha_k\log(|z_1|+\varepsilon_0^{\beta_k})\\&\geq\sum_{k\leq N}\alpha_k\log(|z_1|+\varepsilon_0^{\beta_k})+\sum_{k\geq N+1}\alpha_k\log(|z_1|+\varepsilon_0^{\beta_k})\\&\geq\sum_{k\leq N}\alpha_k\beta_k\log\varepsilon_0+\sum_{k\geq N+1}\alpha_k\log|z_1|\geq 3N\log\varepsilon_0+\sum_{k\geq N+1}\alpha_k\beta_{N+1}\log\varepsilon_0\\&\geq 3N\log\varepsilon_0+2\alpha_{N+1}\beta_{N+1}\log\varepsilon_0=3(N+2)\log\varepsilon_0\end{aligned}$$

and

$$\log(-\log|z_1|)\geq\log(-\beta_N\log\varepsilon_0)=N!\log 2+\log 3+\log(-\log\varepsilon_0).\qquad(*)$$

Set $C_N:=N!\log 2+\log 3+\log(-\log\varepsilon_0)$. Since

$$0<\frac{3(N+2)\log\varepsilon_0}{-C_N}\to 0\ (N\to\infty),$$

it follows from $(*)$ that for large N, we have

$$\frac{3(N+2)\log\varepsilon_0}{-\log(-\log|z_1|)}\leq\frac{3(N+2)\log\varepsilon_0}{-C_N}\leq\frac{\varepsilon_0}{1+\varepsilon}.$$

Hence, for $\varepsilon_0^{\beta_{N+1}}<|z_1|\leq\varepsilon_0^{\beta_N}$, we obtain

$$(1+\varepsilon)\varphi\geq 3(1+\varepsilon)(N+2)\log\varepsilon_0\geq-\varepsilon_0\log(-\log|z_1|)$$

for large enough N. Thus, when $|z_1|$ is small enough, we have

$$\frac{1}{|z_1|^2(-\log|z_1|)^\alpha}e^{-(1+\varepsilon)\varphi}\leq\frac{1}{|z_1|^2(-\log|z_1|)^{\alpha-\varepsilon_0}},$$

which is locally integrable near $(0,z_2,0)\in U$ with $z_2\neq\frac{1}{k},0$ by $\alpha-\varepsilon_0>1$.

Step 3. *Incoherence of the analytic adjoint ideal sheaf* $Adj_D^\alpha(\varphi)$ *near* o.

Suppose that $Adj_D^\alpha(\varphi)$ is a coherent ideal sheaf on U. Then, the zero set $N(Adj_D^\alpha(\varphi))$ of $Adj_D^\alpha(\varphi)$ is an analytic set in U. However, it follows from Step 1 and Step 2 that on U,

$$N(Adj_D^\alpha(\varphi))=\{(0,\frac{1}{k},0)\}\cup\{o\},$$

which is not analytic at o, contradicting to the assumption. □

Remark 2.1. In Theorem 1.1, if the condition $\varphi|_D\not\equiv-\infty$ is not required, then the result also holds for $n=2$ by a slight modification. Indeed, we can take $D=\{z_1=0\}$ and $\varphi(z_1,z_2)=\sum\limits_{k=2}^{\infty}\alpha_k\log(|z_1|+|z_2-\frac{1}{k}|^{\beta_k})$ with $\alpha_k=\frac{1}{2^{k!}}$, $\beta_k=3\cdot 2^{k!}$.

References

[1] H. Guenancia, *Toric plurisubharmonic functions and analytic adjoint ideal sheaves*, Math. Z. 271 (2012), 1011–1035.
[2] D. Kim, *Themes on Non-analytic Singularities of Plurisubharmonic Functions*, Complex Analysis and Geometry, Springer Proceedings in Mathematics & Statistics, vol. 144, Springer, Tokyo, 2015, pp. 197–206.
[3] R. Lazarsfeld, *Positivity in Algebraic Geometry II*, Springer, New York, 2004.
[4] S. Takagi, *Adjoint ideals along closed subvarieties of higher codimension*, J. Reine Angew. Math. 641 (2010), 145–162.

Qi'an Guan: School of Mathematical Sciences, and Beijing International Center for Mathematical Research, Peking University, Beijing, 100871, China.
E-mail address: guanqian@amss.ac.cn

Zhenqian Li: School of Mathematical Sciences, Peking University, Beijing, 100871, China.
E-mail address: lizhenqian@amss.ac.cn